



Genome-wide association study of lung function decline in adults with and without asthma.

Medea Imboden, Emmanuelle Bouzigon, Ivan Curjuric, Adaikalavan Ramasamy, Ashish Kumar, Dana B. Hancock, Jemma B. Wilk, Judith M. Vonk, Gian A. Thun, Valerie Siroux, et al.

► To cite this version:

Medea Imboden, Emmanuelle Bouzigon, Ivan Curjuric, Adaikalavan Ramasamy, Ashish Kumar, et al.. Genome-wide association study of lung function decline in adults with and without asthma.. Journal of Allergy and Clinical Immunology, 2012, 129 (5), pp.1218-28. 10.1016/j.jaci.2012.01.074 . inserm-00744741

HAL Id: inserm-00744741

<https://www.hal.inserm.fr/inserm-00744741>

Submitted on 23 Oct 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genome-wide association study of lung function decline in adults with and without asthma.

Medea Imboden, PhD,^{*a,b}, Emmanuelle Bouzigon, MD, PhD,^{*c,d,e}, Ivan Curjuric, MD,^{a,b},
 Adaikalavan Ramasamy, PhD,^f, Ashish Kumar, MSc,^{a,b,g}, Dana B Hancock, PhD,^{h,i},
 Jemma B Wilk, DSc,^j, Judith M Vonk, PhD,^k, Gian A Thun, MSc,^{a,b}, Valerie Siroux,
 PhD,^{l,m}, Rachel Nadif, PhD,^{n,o}, Florent Monier, MSc,^{c,d,e}, Juan R Gonzalez, PhD,^{p,q},
 Matthias Wjst, MD, MD^r, Joachim Heinrich, PhD,^r, Laura R Loehr, MD, PhD,^s, Nora
 Franceschini, MD, MPH^s, Kari E North, PhD,^t, Janine Altmüller, MD,^u, Gerard H.
 Koppelman, MD, PhD^k, Stefano Guerra, MD, PhD,^{p,q,v,3}, Florian Kronenberg, MD,^w,
 Mark Lathrop, PhD,^{d,x}, Miriam F Moffatt, D.Phil,^y, George T O'Connor, MD, MSc,^{z,1},
 David P Strachan, MD,², Dirkje S Postma, MD, PhD^k, Stephanie J London, MD, DrPH,^h,
 Christian Schindler, PhD,^{a,b}, Manolis Kogevinas, MD,^{p,q,3,4}, Francine Kauffmann, MD,^{n,o},
 Debbie L Jarvis, MD,^f, Florence Demenais, MD,^{c,d,e} and Nicole M Probst-Hensch, PhD,
 PhD,^{#a,b}.

*Contributed equally

corresponding author: N. Probst-Hensch, SwissTPH, Socinstr. 59, 4002 Basel,
 Switzerland. E-Mail: Nicole.Probst@unibas.ch; Telephone: +41 61 284 83 88; Fax +41
 61 284 81 05.

Authors' affiliations:

a –Swiss Tropical and Public Health Institute, Basel, Switzerland

- 24 b – University of Basel, Switzerland
- 25 c – Inserm, UMRS-946, F-75010 Paris, France
- 26 d – Fondation Jean Dausset- Centre d'Etude du Polymorphisme Humain (CEPH), F-
- 27 75010, Paris, France
- 28 e – Univ Paris Diderot, Paris 7, Institut Universitaire d'Hématologie, F-75010, Paris,
- 29 France
- 30 f – Respiratory Epidemiology and Public Health, Imperial College, and MRC-HPA
- 31 Centre for Environment and Health, London, United Kingdom
- 32 g – Wellcome Trust Centre for Human Genetics, University of Oxford, United Kingdom
- 33 h – Epidemiology Branch, Division of Intramural Research, National Institute of
- 34 Environmental Health Sciences, National Institutes of Health, Department of Health and
- 35 Human Services, Research Triangle Park, North Carolina, USA
- 36 i – Behavioral Health Epidemiology Program, Research Triangle Institute International,
- 37 Research Triangle Park, North Carolina, USA
- 38 j – Departments of Neurology and Medicine, Boston University School of Medicine,
- 39 Boston, Massachusetts, USA
- 40 k – Department of Pulmonology, Pediatric Pulmonology and Pediatric Allergy,
- 41 Epidemiology, Beatrix Children's Hospital, Groningen Research Institute for Asthma and
- 42 COPD, University Medical Center Groningen, University of Groningen, The Netherlands
- 43 l – Team of Environmental Epidemiology applied to Reproduction and Respiratory
- 44 Health, Inserm, U823, Grenoble, France
- 45 m – Univ Joseph Fourier, Grenoble, France

46 n – Inserm, U1018, CESP Centre for research in Epidemiology and Population Health,
 47 Respiratory and environmental epidemiology Team, F-94807, Villejuif, France
 48 o – Université Paris Sud, UMRS 1018, F-94807, Villejuif, France
 49 p – Centre for Research in Environmental Epidemiology, Barcelona, Spain
 50 q – CIBER Epidemiologia y Salud Publica, Barcelona, Spain
 51 r – Institute of Epidemiology, Helmholtz Zentrum München, German Research Center
 52 for Environmental Health, Neuherberg, Germany
 53 s – Department of Epidemiology, UNC Gillings School of Global Public Health,
 54 University of North Carolina-Chapel Hill, Chapel Hill, North Carolina, USA
 55 t – Department of Epidemiology and Carolina Center for Genome Sciences, University
 56 of North Carolina-Chapel Hill, Chapel Hill, North Carolina, USA
 57 u – Cologne Center for Genomics (CCG), University of Cologne, Cologne, Germany.
 58 v – Arizona Respiratory Center, University of Arizona, Tucson, AZ, USA
 59 w – Department of Medical Genetics, Molecular and Clinical Pharmacology, Division of
 60 Genetic Epidemiology, Innsbruck Medical University, Austria
 61 x – Commissariat à l'Energie Atomique, Institut de Génomique, Centre National de
 62 Génotypage, Evry, France
 63 y – National Heart and Lung Institute, Imperial College, London, United Kingdom
 64 z – Pulmonary Center, Department of Medicine, Boston University School of Medicine,
 65 Boston, Massachusetts, USA
 66 1 – The National Heart, Lung, and Blood Institute's Framingham Heart Study,
 67 Framingham, Massachusetts, USA

68 2 – Division of Population Health Sciences and Education, St George's, University of
69 London, London, United Kingdom

70 3 – IMIM (Municipal Institute of Medical Research), Barcelona, Spain

71 4 – National School of Public Health, Athens, Greece

72 **Disclosure of potential conflict of interest:**

73

74 Authors declare no conflict of interest.

75

76 **Sources of support:**

77 *Discovery cohorts: ESE (EGEA-SAPALDIA-ECRHS)*

78 **EGEA:** INSERM-Ministry of Research 'Cohortes et Collections' grant (4CH06G). French
79 Ministry of Higher Education and Research, University Paris Diderot-Paris 7, grants
80 from the French Agency for Environmental and Occupational Health Safety (grant
81 AFSSETAPR- SE-2004), the French National Agency for Research (grants ANR 05-
82 SEST-020- 02/05-9-97 and ANR 06-CEBS), PHRC-Paris, Merck Sharp & Dohme
83 (MSD))

84 **SAPALDIA:** Swiss National Science Foundation (grants no 4026-28099,3347CO-
85 108796, 3247BO-104283, 3247BO-104288, 3247BO-104284, 32-65896.01,32-
86 59302.99, 32-52720.97, 32-4253.94); the Federal Office for Forest, Environment and
87 Landscape; the Federal Office of Public Health; the Federal Office of Roads and
88 Transport; the canton's government of Aargau, Basel-Stadt, Basel-Land, Geneva,
89 Luzern, Ticino, Zurich; the Swiss Lung League; the canton's Lung League of Basel
90 Stadt/ Basel Landschaft, Geneva, Ticino and Zurich; Freie Akademische Gesellschaft
91 (FAG); UBS Wealth Foundation.

ECRHS: The co-ordination of ECRHS II was supported by the European Commission,
as part of their Quality of Life programme. The following bodies funded the local studies
in ECRHS II: **Albacete:** Fondo de Investigaciones Santarias (FIS) (grant code:
97/0035-01, 99/0034-01 and 99/0034-02), Hospital Universitario de Albacete,
Consejeria de Sanidad; **Barcelona:** SEPAR, Public Health Service (grant code: R01
HL62633-01), Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01,
99/0034-01 and 99/0034-02) CIRIT (grant code: 1999SGR 00241) Red Respira ISCII;
CIBER Epidemiologia y Salud Pública (CIBERESP), Spain **Basel:** Swiss National
Science Foundation, Swiss Federal Office for Education & Science, Swiss National
Accident Insurance Fund (SUVA), USC NIEHS Center grant 5P30 ES07048; **Bergen:**
Norwegian Research Council, Norwegian Asthma & Allergy Association (NAAF), Glaxo
Wellcome AS, Norway Research Fund; **Erfurt:** GSF-National Research Centre for
Environment & Health, Deutsche Forschungsgemeinschaft (DFG) (grant code FR
1526/1-1); **Galdakao:** Basque Health Dept; **Grenoble:** Programme Hospitalier de
Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health, Direction de la
Recherche Clinique, CHU de Grenoble, Ministere de l'Emploi et de la Solidarite,
Direction Generale de la Sante, Comite des Maladies Respiratoires de l'Isere;
Hamburg: GSF-National Research Centre for Environment & Health, Deutsche
Forschungsgemeinschaft (DFG) (grant code MA 711/4-1); **Ipswich and Norwich:**
Asthma UK (formerly known as National Asthma Campaign); **Huelva:** Fondo de
Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01 and 99/0034-02);
Oviedo: Fondo de Investigaciones Santarias (FIS) (grant code: 97/0035-01, 99/0034-01
and 99/0034-02) ; **Paris:** Ministere de l'Emploi et de la Solidarite, Direction Generale de

115 la Sante, UCB-Pharma (France), Aventis (France), Glaxo France, Programme
116 Hospitalier de Recherche Clinique-DRC de Grenoble 2000 no. 2610, Ministry of Health,
117 Direction de la Recherche Clinique, CHU de Grenoble; **Tartu**: Estonian Science
118 Foundation; **Umeå**: Swedish Heart Lung Foundation, Swedish Foundation for Health
119 Care Sciences & Allergy Research, Swedish Asthma & Allergy Foundation, Swedish
120 Cancer & Allergy Foundation; **Uppsala**: Swedish Heart Lung Foundation, Swedish
121 Foundation for Health Care Sciences & Allergy Research, Swedish Asthma & Allergy
122 Foundation, Swedish Cancer & Allergy Foundation; *Financial support for ECRHS I for*
123 *centres in ECRHS II was provided by:* Ministère de la Santé, Glaxo France, Institut
124 Pneumologique d'Aquitaine, Contrat de Plan Etat-Région Languedoc-Rousillon,
125 CNMATS, CNMRT (90MR/10, 91AF/6), Ministre délégué de la santé, RNSP, France;
126 GSF, and the Bundesminister für Forschung und Technologie, Bonn, Germany;
127 Norwegian Research Council project no. 101422/310; Ministerio Sanidad y Consumo
128 FIS (grants #91/0016060/00E-05E and #93/0393), and grants from Hospital General de
129 Albacete, Hospital General Juan Ramón Jiménez, Consejería de Sanidad Principado
130 de Asturias, Spain; The Swedish Medical Research Council, the Swedish Heart Lung
131 Foundation, the Swedish Association against Asthma and Allergy; Swiss National
132 Science Foundation grant 4026-28099; National Asthma Campaign, British Lung
133 Foundation, Department of Health, South Thames Regional Health Authority, UK. A.R.
134 was supported by the Department of Health, UK and the European Commission as part
135 of GABRIEL contract number 018996 under the Integrated Program LSH-2004-1.2.5-1.
136 **Genotyping of the discovery cohort and part of B58C** was funded by the GABRIEL
137 asthma genetic consortium supported by a contract from the European Commission

(018996) and grants from the French Ministry of Research, the Wellcome Trust (WT084703MA), and Asthma UK.

Replication cohorts:

ARIC: The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C. Infrastructure was partly supported by Grant Number UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research. Work for this manuscript was supported, in part, by the Intramural Research Program of the National Institutes of Health (NIH), National Institute of Environmental Health Sciences (NIEHS, Z01ES043012).

FHS: National Heart, Lung and Blood Institute's Framingham Heart Study (Contract No. N01-HC-25195) and its contract with Affymetrix, Inc for genotyping services (Contract No. N02-HL-6-4278). Dr. Wilk by a Young Clinical Scientist Award from the Flight Attendant Medical Research Institute (FAMRI). A portion of this research utilized the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at Boston University School of Medicine and Boston Medical Center.

B58C: British 1958 Birth Cohort was funded by the Medical Research Council grant G0000934 and the Wellcome Trust grant 068545/Z/02 (<http://www.b58cgene.sgul.ac.uk/>). Genotyping was funded by the Wellcome Trust grant 076113/B/04/Z, by the United States National Institutes of Health and the Juvenile Diabetes Research Foundation U01 DK062418 and by the European Commission Framework Programme 6 (018996).

Dutch Asthma Study: The Dutch Asthma study has been funded by the Netherlands Asthma Foundation grants AF 3.2.07.015; and AF 98.48 and a grant from the University Medical Center Groningen

Word count: Abstract 250 words
Main text 3535 words

ABSTRACT

Background: Genome-wide association studies (GWAS) have identified determinants of chronic obstructive pulmonary disease, asthma and lung function level, however none addressed decline in lung function.

Aim: We conducted the first GWAS on age-related decline in forced expiratory volume in the first second (FEV1) and in its ratio to forced vital capacity (FVC) stratified *a priori* by asthma status.

Methods: Discovery cohorts included adults of European ancestry (1441 asthmatics, 2677 non-asthmatics; Epidemiological Study on the Genetics and Environment of Asthma (EGEA); Swiss Cohort Study on Air Pollution And Lung And Heart Disease In Adults (SAPALDIA); European Community Respiratory Health Survey (ECRHS)). The associations of FEV1 and FEV1/FVC decline with 2.5 million single nucleotide polymorphisms (SNPs) were estimated. Thirty loci were followed-up by *in silico* replication (1160 asthmatics, 10858 non-asthmatics: Atherosclerosis Risk in Communities (ARIC); Framingham Heart Study (FHS); British 1958 Birth Cohort (B58C); Dutch asthma study).

Results: Main signals identified differed between asthmatics and non-asthmatics. None of the SNPs reached genome-wide significance. The association between the height related gene *DLEU7* and FEV1 decline suggested for non-asthmatics in the discovery phase was replicated (discovery $P=4.8 \times 10^{-6}$; replication $P=0.03$) and additional sensitivity analyses point to a relation to growth. The top ranking signal, *TUSC3*, associated with FEV1/FVC decline in asthmatics ($P=5.3 \times 10^{-8}$) did not replicate. SNPs

previously associated with cross-sectional lung function were not prominently associated with decline.

Conclusions: Genetic heterogeneity of lung function may be extensive. Our results suggest that genetic determinants of longitudinal and cross-sectional lung function differ and vary by asthma status.

Key Messages:

- Knowledge regarding genes with pleiotropic effects on asthma, chronic obstructive pulmonary disease as well as on lung function level and its longitudinal course is limited.
- This first GWAS meta-analysis on lung function decline conducted separately in non-asthmatic and asthmatic cohort participants suggests that genetic determinants of lung function decline are different in the two groups.
- The results further suggest that previously identified genetic determinants of cross-sectional lung function are not major determinants of the decline.

Capsule summary:

This meta-analysis provides evidence for genetic heterogeneity of lung function between asthmatics and non-asthmatics; and between cross-sectionally and longitudinally measured lung function. The study adds evidence for the role of height-related genes in lung health.

218

219 This article has in support of the manuscript online repository materials.

220

221 **Keywords:**

222 Asthma, cohort studies, genome-wide association, lung function decline, heterogeneity

223

224 **Abbreviations:**

225 ARIC, Atherosclerosis Risk in Communities Study

226 ATS, American Thoracic Society

227 B58C, British 1958 Birth Cohort

228 chr, chromosome

229 COPD, chronic obstructive pulmonary disease

230 ECRHS, European Community Respiratory Health Survey

231 EGEA, Genetics and Environment of Asthma

232 FEV1, forced expiratory volume in the first second

233 FHS, Framingham Heart Study

234 FVC, forced vital capacity

235 GWAS, genome-wide association studies

236 HapMap, Haplotype Map Project

237 Q-Q, Quantile-quantile

238 SAPALDIA, Swiss Cohort Study on Air Pollution And Lung And Heart Disease In Adults

239 SNP, single nucleotide polymorphism

INTRODUCTION

Low lung function is a feature of both asthma and chronic obstructive pulmonary disease (COPD), with twin studies demonstrating strong heritability (0.51 to 0.77) for forced expiratory volume in the first second (FEV1)^{1, 2}. The two respiratory diseases and lung function itself share predisposing and phenotypic features, including increased airway responsiveness and atopy as well as exogenous risk factors^{3, 4}. Genome-wide association studies (GWAS) have identified novel genetic loci for asthma⁵⁻¹⁰, COPD¹¹⁻¹⁴, and lung function¹⁵⁻¹⁸ and provide the opportunity to study agnostically their overlap in genetic background¹⁹. Some of the implicated genes, such as *PDE4D*, support a link between asthma and COPD which may be rooted in shared pathways during lung development²⁰. However, the majority of the genes implicated in asthma or COPD GWAS analyses have not been identified as top association signals in GWAS for lung function in the general population¹⁵⁻¹⁸, with the exception of *HHIP* and *FAM13A* being associated with both lung function¹⁵⁻¹⁸ and COPD¹¹⁻¹⁴. Several lines of evidence suggest that different genes influence lung function in asthmatics and in non-asthmatics. Genome-scans in family based linkage studies identified some, but overall limited overlap between chromosomal regions linked to lung function in asthmatics²¹, COPD patients²² and in the general population²³ and it has been suggested that genetic variation may be more important for lung function in asthma after adjusting for smoking and body size differences^{21, 24, 25}.

Here, we present results from the first lung function GWAS conducted separately for asthmatics and non-asthmatics. This current study also focuses on the rate of lung function decline in adults instead of cross-sectional lung function parameters tested in

263 previous GWAS¹⁵⁻¹⁸. The discovery cohorts included two population-based studies
264 (SAPALDIA and ECRHS) and one asthma family-based study (EGEA), all of European
265 ancestry with highly comparable and standardized assessment of respiratory health
266 parameters including spirometry from two time points ten years apart. These three
267 studies had been included in the GWAS for asthma conducted by the GABRIEL
268 consortium⁷. Replication cohorts included three population-based cohorts (FHS, ARIC,
269 B58C) and one family-based asthma study (the Dutch Asthma Study).

METHODS

- Discovery cohorts and study population: Three large multi-centric cohorts EGEA²⁶, SAPALDIA²⁷ and ECRHS²⁸ constitute the ESE-consortium. Personal factors of relevance to lung function decline were assessed by interview and anthropometric measurements at baseline and follow-up. Participants included in discovery phase were derived from the nested asthma case/control samples (SAPALDIA and ECRHS) or from the entire study population (EGEA) subjected to genome-wide genotyping in the context of the GABRIEL asthma GWAS⁷. Baseline and follow-up examination were roughly 10 years apart. The analysis was restricted to adult participants (age ≥ 18 years at the time of the baseline spirometry) with complete information on age, height and sex as well as valid lung function measure from both surveys. Cohort study protocols were in agreement with the Declaration of Helsinki and obtained ethical approval from their respective regional and/or national review boards.

- Lung function assessments, asthma status and genotypes: At each visit, a minimum of two acceptable forced expiratory flows, forced vital capacity (FVC) and forced expiratory volume in the first second (FEV1) complying with American Thoracic Society criteria were obtained²⁶⁻²⁹. No bronchodilator was administered. Based on questionnaire data, asthmatics were defined as asthma self-report at any of the completed surveys and family-based studies considered additional clinical asthma criteria (see online repository). Genotyping for discovery cohorts was centrally performed on the Illumina Human 610quad BeadChip at the Centre National de Génomique (CNG, Evry, France)⁷. Imputation of genotypes based on Hapmap2 reference panel, investigation of

population stratification and quality control criteria are described in Figure E1 and Table E1 in the Online Repository.

-Replication Cohorts: Four cohorts of European ancestry with available genome-wide data, ARIC³⁰, FHS¹⁵; B58C³¹; Dutch asthma study³² were used for replication. Subjects included in the current analysis were older than 24 years, had complete information on covariates (age, height, and sex) and valid lung function measures from at least two time-points. The lung function measurements were conducted at least ten years apart, except three years apart for ARIC (Table I). Distinct genotype data platforms and imputation software were used (Table E11, Online Repository).

- Statistical analysis: Annual decline in FEV1 and FEV1/FVC was calculated as difference between follow-up and baseline spirometric measurements (mL for FEV1 and % for FEV1/FVC) divided by the duration of follow-up in years. Standardized residuals were derived from sex-specific linear regression models adjusted for age, height and study centre in asthmatics and non-asthmatics separately. Comparability between studies of standardized residuals was tested using Wilcoxon-Mann-Whitney test (P>0.94). The standardized residuals were used as dependent variable and regressed on genome-wide single nucleotide polymorphisms (SNPs) adjusted for study-specific principal components capturing population ancestry (see online supplement for details). Study-specific SNP effect estimates were combined through meta-analysis using fixed and random effects models. We used a threshold of $P < 5 \times 10^{-8}$ (the Bonferroni adjustment for one million independent tests) to declare a pooled effect as genome-wide significant. Selection criteria for replication loci are described in the methods section of the online repository. SNPs with suggestive evidence of association with

decline in FEV1 or FEV1/FVC were chosen for *in silico* replication (Table EIII, Online Repository). Study-specific regression models and meta-analyses across replication cohorts were as described for the discovery phase. Replication cohorts with spirometry data from more than two different time points modelled the lung function decline phenotype by fitting a least-squares slope using the available data (FHS, Dutch asthma study). $P \leq 0.05$ was considered as statistically significant at the replication level.

The results of the main meta-analyses for the top 1000 SNPs are available in the online repository (Table EIV A to D, Online Repository). We also conducted a meta-analysis by combining non-asthmatic and asthmatic samples and tested for heterogeneity between these samples (Table EV, Online Repository). Additional sensitivity analyses were done by: a) restricting the GWAS sample to subjects aged 30 and older for FEV1 decline (Table EIV E and F, Online Repository); b) conducting GWAS analyses on percent change instead of absolute annual decline in lung function (Table EIV G to J, Online Repository); c) investigating smoking stratified joint effects for replications SNPs (Table EVI, Online Repository); d) excluding ARIC, a cohort having substantially shorter follow-up time than the other cohorts (three years instead of ten years) from replication analyses (Table EVII, Online Repository). Methods and results of these additional analyses are described in the online repository.

RESULTS

Characteristics of the study populations

The cohorts included in this study differed by age and type of recruitment, and accordingly in lung function and the proportion of subjects with FEV1/FVC below 70% (Table I, Table EVIII, Online Repository). Baseline lung function parameters, but not their annual changes were lower in asthmatics when compared to non-asthmatics in each study. The proportion of never smokers was comparable among asthmatics, but varied among non-asthmatics (ranging from 28.5% in B58C to 46.5% in EGEA). No substantial differences in the smoking prevalence between people with and without asthma were observed within each study. Comparing the discovery cohorts in more detail (Table EVIII, Online Repository), atopy (total IgE ≥ 100 kU/ml) and hay fever were more prevalent in both asthmatics and non-asthmatics from EGEA when compared to ECRHS and SAPALDIA. Current asthma was more prevalent (84.4%) in EGEA than in SAPALDIA (25.5%) or ECRHS (43.3%) and the prevalence of a positive family history for asthma was also highest in EGEA, in agreement with the study design. Asthmatics from EGEA had a younger age of disease onset due to the mode of recruitment of the proband.

Main findings from meta-analyses of discovery and replication phase

In the discovery phase, GWAS meta-analysis of decline in FEV1 and FEV1/FVC was conducted in 2677 non-asthmatics and in 1441 asthmatics. Genomic inflation factors were low for both lung function parameters ($\lambda < 1.047$, Table EIX, Online Repository) suggesting minimal unaccounted population stratification. The replication panel included

a total of 10'858 non-asthmatics and 1'138 asthmatics. Thirty lead SNPs belonging to 30 loci ($5 \times 10^{-8} < P_{\text{discovery}} < 6 \times 10^{-5}$) were chosen for replication.

The four lung function parameter- and asthma-specific meta-analyses identified one association signal that almost reached the genome-wide significance level ($P = 5.3 \times 10^{-8}$) at the locus 8p22 containing the *TUSC3* gene for FEV1/FVC decline in asthmatics while all other signals had $P < 5 \times 10^{-7}$ (Figure I), but this signal was not associated with FEV1/FVC decline in asthmatics in the replication sample. The only locus of the selected replication candidate loci that formally replicated was 13q14.3, containing the *DLEU7* gene, associated with decline in FEV1 in the non-asthmatics ($P_{\text{discovery}} = 4.8 \times 10^{-6}$ and $P_{\text{replication}} = 0.03$).

In the global *post hoc* analysis combining both asthmatics and non-asthmatics (N=4118), a striking finding was the absence of any pronounced association signals ($P > 1 \times 10^{-6}$) despite increased statistical power. This was in agreement with the minimal overlap of association signals observed in asthmatics and non-asthmatics separately. Most signals at $P < 10^{-5}$ from the asthma-stratified analysis in the discovery phase exhibited statistically significant heterogeneity of effects between the two groups (Table II). At the replication stage, none of the replication SNPs was associated with lung function decline in asthmatics and non-asthmatics combined.

Association signals for annual decline in FEV1 in non-asthmatics

Of fifteen SNPs associated at $P < 10^{-5}$ with decline in FEV1 in non-asthmatics ten were clustered at position 112.3 Mb on chromosome 9, containing genes *TXN*, *MUSK* and *SVEP1*. Two of the 15 SNPs were located at 13q14.3 in a locus containing the *DLEU7*

gene; three SNPs belonged to three distinct loci. The association of lead and proxy SNPs in *DLEU7* (Figure II), but not *TXN/MUSK/SVEP1* (Figure EII) or the other SNPs (Table II) replicated. The G-allele of SNP rs9316500 near the *DLEU7* gene was positively associated with annual FEV1 decline in the discovery cohorts ($P=4.8 \times 10^{-6}$) and in the replication cohorts ($P=0.026$). Although heterogeneity between studies was not significant ($P=0.61$), the combined P value did not reach the genome-wide level ($P=5.7 \times 10^{-5}$).

Association signals for annual decline in FEV1 in asthmatics

Eighteen SNPs in nine distinct chromosomal locations were associated with decline in FEV1 in asthmatics at $P < 10^{-5}$. None of the loci selected for *in silico* replication was confirmed (Table II).

Association signals for annual decline in FEV1/FVC in non-asthmatics

Seven loci showed association with FEV1/FVC decline in non-asthmatics at $10^{-6} < P < 10^{-5}$, but no locus selected for replication was confirmed (Table II).

Association signals for annual decline in FEV1/FVC in asthmatics

Twelve SNPs at the locus 8p22 containing the gene *TUSC3* at 15.68Mb were associated with FEV1/FVC decline at $P < 10^{-7}$ in asthmatics (Figure I). Regional locus plot and forest plot are presented in the online repository (Figure EIII). The top association signals in this locus were conferred by distinct SNPs in each cohort, though apparently they were located in the same putative haplotype segment in SAPALDIA and

in EGEA (Figure EIV, Online Repository). There was no statistically significant association in ECRHS. Meta-analysis of the discovery samples identified SNP rs4831760 as top signal in *TUSC3* gene, but heterogeneity between discovery studies was borderline significant ($P=0.07$). The C-allele ($P=5.3 \times 10^{-8}$) was positively associated with annual decline in FEV1/FVC in asthmatics (Beta=0.22 \pm 0.04 (standard error); Table II). However this association was not replicated ($P=0.80$). In the meta-analysis combining discovery and replication samples the resulting P-value for rs4831760 was 2.8×10^{-5} . All but the Dutch asthma study, exhibited effect estimates in the same direction as the discovery panel. Two other candidate loci (*MPP7* and *SYNE2*) also failed replication testing.

SNPs previously associated in GWAS meta-analyses on cross-sectional lung function

The associations of top hit SNPs from previous GWAS meta-analyses on cross-sectional lung function^{11, 15-18} and a replication study in asthmatics³³ were assessed separately for asthmatics and non-asthmatics in the discovery cohorts. Associations were assessed for both, lung function parameters of decline (annual decline and percent change) and cross-sectional lung function level. Overall, a subset of variants and loci showed replication of association with cross-sectional lung function in either non-asthmatics or asthmatics. Few of the loci showed strong association with decline in lung function. We present associations at $P < 0.05$ in Table III and those at $P \geq 0.05$ in Table EX in the online repository.

For baseline FEV1, we observed associations for SNPs belonging to 4q24 (*GSTCD*, rs11731417, $P=1.3 \times 10^{-4}$) and 15q23 (*THSD4*, rs1913768, $P=0.003$). Associations with

baseline FEV1 were mainly restricted to non-asthmatics. For baseline FEV1/FVC, associations of SNPs of *THSD4* were prominent (e.g. rs12899618, $P=3.3 \times 10^{-4}$) and again restricted to non-asthmatics.

For decline phenotypes of FEV1, we observed associations for SNPs in regions 6p21 (*DAAM2*, $0.003 < P < 0.02$) and 4q28 (*HHIP*, $0.02 < P < 0.05$) among asthmatics and in *THSD4* ($0.003 < P < 0.04$) among non-asthmatics. The strongest associations observed for decline phenotypes of FEV1/FVC were two SNPs in *MMP15* (16q13, $0.003 < P < 0.002$) in non-asthmatics, only. Association in the combined sample of asthmatics and non-asthmatics did not substantially alter the results.

Summary of findings from sensitivity analyses

We observed in non-asthmatics, aged 30 years and more, that *MUSK* and *DLEU7* were no longer prominently associated with FEV1 decline, but SNPs in other genes remained strongly associated (*ZIC1*, rs6785065, $P=2.3 \times 10^{-5}$; *UBL3*, rs278037, $P=4.8 \times 10^{-5}$).

Results of the GWAS on percent change in lung function showed that the FEV1 association signal for *DLEU7* in the non-asthmatics was no longer significant; however the signals for *MUSK* (rs1889321, $P=2.92 \times 10^{-7}$) and other loci remained unaltered (*ZIC1*, rs6785065, $P=2.0 \times 10^{-5}$; *KIRREL3*, rs11604082, $P=4.1 \times 10^{-6}$; *KIAA2117*, rs10082549, $P=2.7 \times 10^{-6}$). Top signals associated with decline in FEV1/FVC in asthmatics remained unaltered for *TUSC3* (rs4831760, $P=5.2 \times 10^{-8}$) and for *SYNE2* (rs7144584, $P=6.4 \times 10^{-7}$) after taking baseline lung function into account.

Smoking stratified analyses of the replication SNPs revealed no substantial difference in association between ever and never smokers except for a few SNPs belonging to loci containing *SYNE2*, *RORA*, *BCAS1*, or *PLXNA4* genes.

Replication meta-analysis excluding the ARIC data substantially reduced sample size in non-asthmatics and the association of *DLEU7* with decline of FEV1 was no longer significant. Instead two loci for association with decline in FEV1 in asthmatics (*PLXNA4*, rs10808265, $P_{\text{discovery}}=1.7 \times 10^{-6}$, $P_{\text{replication}}=0.02$ and *SLC45A3*, rs16856186, $P_{\text{discovery}}=8.9 \times 10^{-6}$, $P_{\text{replication}}=0.04$) and one locus, FLJ25393, for decline in FEV1/FVC in non-asthmatics (rs2658782, $P_{\text{discovery}}=4.3 \times 10^{-6}$, $P_{\text{replication}}=0.03$) gained statistical significance.

DISCUSSION

A main result of this study is the observed genetic heterogeneity of lung function decline between asthmatics and non-asthmatics. When we combined the two groups in the discovery phase we observed no genome-wide significant association signal despite larger sample size. All top hit association signals detected by the asthma stratified analysis showed significant heterogeneity according to the disease status. In the replication phase, this heterogeneity was also confirmed for the *DLEU7* locus which was associated with FEV1 decline in non-asthmatics only. Finally, many of the SNPs identified by previous GWAS on lung function exhibited associations specific to asthma status.

The finding of genetic heterogeneity in lung function reported here is consistent with available evidence. Differences in familial segregation of FEV1 in asthmatic and non-asthmatic families previously suggested genetic heterogeneity between these two groups²⁴. Agnostic studies investigating genetic determinants of lung function in both, family-based^{21, 22, 34-37} and population-based samples^{15-18, 23, 25} produced little overlap in chromosomal regions. Genome-wide scans on lung function in asthma^{21, 38} or COPD²² families also suggested a heterogeneous genetic architecture of lung function.

Nevertheless, some previously reported overlapping linkage regions for the ratio of FEV1 over vital capacity (FEV1/VC) and FEV1 over the forced vital capacity (FEV1/FVC) in families with asthma and COPD^{21, 22} suggest that at least some gene(s) could be important in the development of airway obstruction in both diseases.

Furthermore, genetic polymorphisms in glutathione S-transferases³⁹⁻⁴² as well as *ADAM-33*⁴³⁻⁴⁶ were associated with lower lung function at all ages and in different subgroups of the population (general population, patients with COPD and asthma). Gene-lung function associations that are of relevance to several population and patient strata may be determined specifically by complex gene-gene and gene-environment interactions, as suggested for lung function decline and its complex association with estrogen receptor 1 polymorphisms, smoking, steroid use, and gender^{32, 47}. While ignored in ours as well as previous GWAS, such effect modifications should be considered in the future⁴⁸.

Results from the Busselton Health Study on familial aggregation and heritability of adult lung function previously suggested the existence of genetic determinants of adult lung function independent of asthma, atopy, cigarette smoking, height, age or sex²⁵. Consistent with these results, neither asthma, atopy and COPD genes previously identified in large GWAS^{5-9, 11} nor genes related to smoking behavior⁴⁹ were associated with lung function decline in our study. The association of FEV1 decline with a gene related to height, *DLEU7*, was ranking high, but only in subjects without asthma (rs9316500, $P_{\text{discovery}}=4.8 \times 10^{-6}$; $P_{\text{replication}}=0.03$). *DLEU7* gene product and expression remain poorly characterized, but its mRNA has been detected in the lung. The *DLEU7* locus was identified as a determinant of adult height in previous GWAS meta-analyses⁵⁰⁻⁵². Three other height genes, *HHIP*, *GPR126* and *PTCH*, were associated with cross-sectional lung function¹⁵⁻¹⁷. All of these lung function models including ours were adjusted for adult height. The observed association, related to both *HHIP* and

504 *DLEU7* being associated with peak height velocity in infancy⁵¹, suggests that aspects
505 beyond adult height influence lung function and possibly its response to non-genetic
506 determinants. Several genes implicated in respiratory diseases indicate that early lung
507 development impacts respiratory health later in life²⁰. Sensitivity analyses are supportive
508 for a growth-specific role of *DLEU7*. The association of genetic variants in *DLEU7* with
509 decline in FEV1 disappeared in analyses considering baseline lung function or restricted
510 to subjects above age 30 with no remaining physiologic lung growth. There might be a
511 link between physiologic growth and unregulated cell differentiation as the *DLEU7* gene
512 is also a proposed tumor suppressor gene in chronic lymphocytic leukemia⁵³⁻⁵⁵.
513 Evidence emerges for a role of *DLEU7* in counterbalancing the proliferative impact of
514 NF- κ B on various cell types⁵⁶. The potential role of the gene product of *TUSC3*, a
515 proposed tumor suppressor gene⁵⁷, in lung physiology is discussed in the Online
516 Repository.

517 None of the SNPs identified in GWAS of cross-sectional lung function¹⁵⁻¹⁸ ranked high in
518 this current GWAS on lung function decline. A strong risk factor for accelerated lung
519 function decline in adulthood is cigarette smoking, but our study was too small to assess
520 gene smoking interaction at the GWAS level. We had decided *a priori* against smoking
521 adjustment as it is not a confounder, and any link between genotype and smoking is
522 likely to be, at least in part, in the same causal pathway (e.g. gene products
523 metabolizing tobacco constituents or influencing smoking behavior). Their identification
524 as determinants of lung function decline is of public health importance. Consistent with
525 previous GWAS on cross-sectional lung function¹⁵⁻¹⁸, neither the *TUSC3* (heterogeneity
526 between ever/never smokers $P=0.98$) nor other top hit signals were modified by

smoking except for SNPs in *SYNE2*, *RORA*, *BCAS1* and *PLXN4*. Arguments for biologic plausibility are mentioned in the Online Repository.

The strength of the present study is the longitudinal design of all cohorts included.

Repeated spirometric assessments within the same subject is thought to capture more

precisely exogenous factors and genes leading to accelerated loss of lung function in

adulthood⁵⁸. The discovery cohorts shared comparable questionnaire and spirometry

protocols and they were specifically designed to investigate environmental and genetic

causes of lung function decline and asthma in a standardized way. Each study has two

measures of pre-bronchodilator lung function about ten years apart, but clearly our

findings would be more robust if further lung function measures were available over an

even longer period of follow-up. All discovery cohorts have used the same genotyping

platform and stringent quality control criteria have been applied.

Sample size is a limitation of this study, and remains a general challenge in lung

function studies with a need for high phenotypic comparability as spirometry results are

sensitive to technicians and devices used⁵⁹. The pre-bronchodilation lung function

measurements in our and previous lung function GWAS do not allow to differentiate

reversible from non-reversible obstruction to airflow. Populations included in this study

differed by age which is also reflected by the diverging proportion of subjects with

FEV1/FVC <0.7 at follow-up between the discovery cohorts. Discovery and replication

populations also differ by time spacing between the spirometry assessments. We can

only speculate of on the overall impact of such differences. We do note that replication

results were sensitive to the exclusion of ARIC data (the study with highest mean age,

largest annual decline, and shortest follow-up time).

Other limitations are shared with any GWAS meta-analyses investigating complex phenotypes such as lack in power for investigating gene-environment interactions or studying subgroups of diseases. As the sample size of our study was comparatively small, especially for the asthmatic sample in the replication phase, we had limited ability to address differences in asthma sub-phenotypes or the impact of asthma medication intake. It is also likely that a substantial part of complex disease may be explained by rare mutations not considered by current GWAS. Finally, assessing the joint effect of SNPs having small effects individually and potentially interacting with each other remains another challenge.

In conclusion, this first GWAS meta-analysis on lung function decline provides suggestive evidence for genetic heterogeneity between persons with and without asthma and between cross-sectionally and longitudinally measured lung function. Consistent with cross-sectional GWAS, our results are also suggestive of height related genes playing a role. Further studies in this area would be enhanced by greater comparability of age range, spacing of lung function assessments, and asthma sub-phenotypes (including treatment) to decrease phenotypic heterogeneity and therefore increase statistical power to detect true association candidate loci⁶⁰.

ACKNOWLEDGMENTS

EGEA: We thank the EGEA cooperative group: *Coordination:* F Kauffmann; F Demenais (genetics); I Pin (clinical aspects). *Respiratory epidemiology* : Inserm U 700, Paris M Korobaeff (Egea1), F Neukirch (Egea1); Inserm U707, Paris : I Annesi-Maesano ; Inserm CESP/U 1018, Villejuif : F Kauffmann, N LeMoual, R Nadif, MP Oryszczyn ; Inserm U 823, Grenoble : V Siroux *Genetics* : Inserm U 393, Paris : J Feingold ; Inserm U 946, Paris : E Bouzigon , F Demenais, MH Dizier ; CNG, Evry : I Gut , M Lathrop. Clinical centers : Grenoble : I Pin, C Pison; Lyon : D Ecochard (Egea1), F Gormand, Y Pacheco ; Marseille : D Charpin (Egea1), D Vervloet ; Montpellier : J Bousquet ; Paris Cochin : A Lockhart (Egea1), R Matran (now in Lille) ; Paris Necker : E Paty, P Scheinmann ; Paris-Trousseau : A Grimfeld, J Just. *Data and quality management* : Inserm ex-U155 (Egea1) : J Hochez ; Inserm CESP/U 1018, Villejuif : N Le Moual, Inserm ex-U780 : C Ravault ; Inserm ex-U794 : N Chateigner ; Grenoble : J Ferran. The authors thank all those who participated to the setting of the study and on the various aspects of the examinations involved: interviewers, technicians for lung function testing, coders, those involved in quality control, data management and all those who supervised the study in all centers. The authors are grateful to the three CIC-Inserm of Necker, Grenoble and Marseille who supported the study and in which subjects were examined. They are indebted to all the individuals who participated without whom that study would not have been possible. **SAPALDIA:** The study could not have been done without the help of the study participants, technical and administrative support and the medical teams and field workers at the local study sites. Local fieldworkers : Aarau: M Broglie, M Bünter, D Gashi, Basel: R Armbruster, T

591 Damm, U Egermann, M Gut, L Maier, A Vögelin, L Walter, Davos: D Jud, N Lutz,
 592 Geneva: M Ares, M Bennour, B Galobardes, E Namer, Lugano: B Baumberger, S
 593 Boccia Soldati, E Gehrig-Van Essen, S Ronchetto, Montana: C Bonvin, C Burrus,
 594 Payerne: S Blanc, AV Ebinger, ML Fragnière, J Jordan, Wald: R Gimmi, N Kourkoulos,
 595 U Schafroth. Administrative staff: N Bauer, D Baehler, C Gabriel, R Gutknecht.
 596 SAPALDIA Team: *Study directorate*: T Rochat, , JM Gaspoz, N Künzli, LJS Liu, NM
 597 Probst Hensch, C Schindler. *Scientific team*: JC Barthélémy, W Berger, R Bettschart, A
 598 Bircher, G Bolognini, O Brändli, C Brombach, M Brutsche, L Burdet, M Frey, U Frey,
 599 MW Gerbase, D Gold, E de Groot, W Karrer, R Keller, B Knöpfli, B Martin, D Miedinger,
 600 U Neu, L Nicod, M Pons, F Roche, T Rothe, E Russi, P Schmid-Grendelmeyer, A
 601 Schmidt-Trucksäss, A Turk, J Schwartz, D. Stolz, P Straehl, JM Tschopp, A von
 602 Eckardstein, E Zemp Stutz. *Scientific team at coordinating centers*: M Adam, E Boes,
 603 PO Bridevaux, D Carballo, E Corradi, I Curjuric, J Dratva, A Di Pasquale, L Grize, D
 604 Keidel, S Kriemler, A Kumar, M Imboden, N Maire, A Mehta, F Meier, H Phuleria, E
 605 Schaffner, GA Thun, A Ineichen, M Ragettli, M Ritter, T Schikowski, G Stern, M
 606 Tarantino, M Tsai, M Wanner. **ECRHS**: The European Community Respiratory Health
 607 Survey is a collaboration of European research groups many of whom also agreed to
 608 provide blood samples for genotyping as part of the GABRIEL initiative. Investigators in
 609 the collaborating centres are Debbie Jarvis, Matthias Wjst, ,Manolis Kogevinas, Rain
 610 Jogi, Christer Janson, Karl Franklin, Ernst Omenaas, Benedicte Leynaert, Isabelle Pin,
 611 Joachim Heinrich, Nino Kuenzli, Nicole M. Probst-Hensch, Josep M. Anto, Jordi Sunyer,
 612 Jose-Antonio Maldonado, Jesus Martinez-Moratalla, Isabel Urritia, Felix Payo. **EGEA**,
 613 **SAPPALDIA and ECRHS** were part of the **GABRIEL** Consortium, a European 6th

614 Framework Research project on asthma genetics, which allowed us to obtain the
615 genotype information used in this analysis. **ARIC:** The authors thank the staff and
616 participants of the ARIC study for their important contributions. Grace Chiu at Westat
617 Inc. (Research Triangle Park, NC), Shuangshuang Dai at the National Institute of
618 Environmental Health Sciences, and Richard Howard at the University of North Carolina
619 School of Public Health provided data management and programming assistance. **FHS**
620 research was conducted using data and resources from the Framingham Heart Study of
621 the National Heart, Lung, and Blood Institute of the National Institutes of Health and
622 Boston University School of Medicine. The analyses reflect intellectual input and
623 resource development from the Framingham Heart Study investigators participating in
624 the SNP Health Association Resource (SHARe) project. **B58C:** We acknowledge use of
625 phenotype and genotype data from the British 1958 Birth Cohort DNA collection, funded
626 by the Medical Research Council grant G0000934 and the Wellcome Trust grant
627 068545/Z/02. (<http://www.b58cgene.sgul.ac.uk/>). Genotyping for the B58C-WTCCC
628 subset was funded by the Wellcome Trust grant 076113/B/04/Z. The B58C-T1DGC
629 genotyping utilized resources provided by the Type 1 Diabetes Genetics Consortium, a
630 collaborative clinical study sponsored by the National Institute of Diabetes and Digestive
631 and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases
632 (NIAID), National Human Genome Research Institute (NHGRI), National Institute of
633 Child Health and Human Development (NICHD), and Juvenile Diabetes Research
634 Foundation International (JDRF) and supported by U01 DK062418. B58C-T1DGC
635 GWAS data were deposited by the Diabetes and Inflammation Laboratory, Cambridge
636 Institute for Medical Research (CIMR), University of Cambridge, which is funded by

637 Juvenile Diabetes Research Foundation International, the Wellcome Trust and the
638 National Institute for Health Research Cambridge Biomedical Research Centre; the
639 CIMR is in receipt of a Wellcome Trust Strategic Award (079895). The B58C-GABRIEL
640 genotyping was supported by a contract from the European Commission Framework
641 Programme 6 (018996) and grants from the French Ministry of Research.

REFERENCES

1. Hankins D, Drage C, Zamel N, Kronenberg R. Pulmonary function in identical twins raised apart. *Am Rev Respir Dis* 1982; 125:119-21.
2. Redline S, Tishler PV, Rosner B, Lewitter FI, Vandenberg M, Weiss ST, et al. Genotypic and phenotypic similarities in pulmonary function among family members of adult monozygotic and dizygotic twins. *Am J Epidemiol* 1989; 129:827-36.
3. Guerra S. Asthma and chronic obstructive pulmonary disease. *Curr Opin Allergy Clin Immunol* 2009; 9:409-16.
4. Postma DS, Boezen HM. Rationale for the Dutch hypothesis. Allergy and airway hyperresponsiveness as genetic factors and their interaction with environment in the development of asthma and COPD. *Chest* 2004; 126:96S-104S; discussion 59S-61S.
5. Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* 2007; 448:470-3.
6. Himes BE, Hunninghake GM, Baurley JW, Rafaels NM, Sleiman P, Strachan DP, et al. Genome-wide association analysis identifies PDE4D as an asthma-susceptibility gene. *Am J Hum Genet* 2009; 84:581-93.
7. Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med* 2010; 363:1211-21.
8. Sleiman PM, Flory J, Imielinski M, Bradfield JP, Annaiah K, Willis-Owen SA, et al. Variants of DENND1B associated with asthma in children. *N Engl J Med* 2010; 362:36-44.
9. DeWan AT, Triche EW, Xu X, Hsu LI, Zhao C, Belanger K, et al. PDE11A associations with asthma: results of a genome-wide association scan. *J Allergy Clin Immunol* 2010; 126:871-3 e9.
10. Hancock DB, Romieu I, Shi M, Sienra-Monge JJ, Wu H, Chiu GY, et al. Genome-wide association study implicates chromosome 9q21.31 as a susceptibility locus for asthma in mexican children. *PLoS Genet* 2009; 5:e1000623.
11. Pillai SG, Ge D, Zhu G, Kong X, Shianna KV, Need AC, et al. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet* 2009; 5:e1000421.
12. Pillai SG, Kong X, Edwards LD, Cho M, Anderson WH, Coxson HO, et al. Loci Identified by Genome-wide Association Studies Influence Different Disease-related Phenotypes in COPD. *Am J Respir Crit Care Med* 2010.
13. Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Hersh CP, et al. Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet* 2010; 42:200-2.
14. Kong X, Cho MH, Anderson W, Coxson HO, Muller N, Washko G, et al. Genome-wide Association Study Identifies BICD1 as a Susceptibility Gene for Emphysema. *Am J Respir Crit Care Med* 2011; 183:43-9.
15. Wilk JB, Chen TH, Gottlieb DJ, Walter RE, Nagle MW, Brandler BJ, et al. A genome-wide association study of pulmonary function measures in the Framingham Heart Study. *PLoS Genet* 2009; 5:e1000429.

16. Hancock DB, Eijgelsheim M, Wilk JB, Gharib SA, Loehr LR, Marcianti KD, et al. Meta-analyses of genome-wide association studies identify multiple loci associated with pulmonary function. *Nat Genet* 2010; 42:45-52.
17. Repapi E, Sayers I, Wain LV, Burton PR, Johnson T, Obeidat M, et al. Genome-wide association study identifies five loci associated with lung function. *Nat Genet* 2010; 42:36-44.
18. Artigas MS, Loth DW, Wain LV, Gharib SA, Obeidat M, Tang W, et al. Genome-wide association and large-scale follow up identifies 16 new loci influencing lung function. *Nat Genet* 2011.
19. Weiss ST. Lung function and airway diseases. *Nat Genet* 2010; 42:14-6.
20. Weiss ST. What genes tell us about the pathogenesis of asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2010; 181:1170-3.
21. Postma DS, Meyers DA, Jongepier H, Howard TD, Koppelman GH, Bleecker ER. Genomewide screen for pulmonary function in 200 families ascertained for asthma. *Am J Respir Crit Care Med* 2005; 172:446-52.
22. Silverman EK, Palmer LJ, Mosley JD, Barth M, Senter JM, Brown A, et al. Genomewide linkage analysis of quantitative spirometric phenotypes in severe early-onset chronic obstructive pulmonary disease. *Am J Hum Genet* 2002; 70:1229-39.
23. Wilk JB, DeStefano AL, Arnett DK, Rich SS, Djousse L, Crapo RO, et al. A genome-wide scan of pulmonary function measures in the National Heart, Lung, and Blood Institute Family Heart Study. *Am J Respir Crit Care Med* 2003; 167:1528-33.
24. Holberg CJ, Morgan WJ, Wright AL, Martinez FD. Differences in familial segregation of FEV1 between asthmatic and nonasthmatic families. Role of a maternal component. *Am J Respir Crit Care Med* 1998; 158:162-9.
25. Palmer LJ, Knuiman MW, Divitini ML, Burton PR, James AL, Bartholomew HC, et al. Familial aggregation and heritability of adult lung function: results from the Busselton Health Study. *Eur Respir J* 2001; 17:696-702.
26. Kauffmann F, Dizier MH, Pin I, Paty E, Gormand F, Vervloet D, et al. Epidemiological study of the genetics and environment of asthma, bronchial hyperresponsiveness, and atopy: phenotype issues. *Am J Respir Crit Care Med* 1997; 156:S123-9.
27. Downs SH, Schindler C, Liu LJ, Keidel D, Bayer-Oglesby L, Brutsche MH, et al. Reduced exposure to PM10 and attenuated age-related decline in lung function. *N Engl J Med* 2007; 357:2338-47.
28. The European Community Respiratory Health Survey II. *Eur Respir J* 2002; 20:1071-9.
29. Standardization of Spirometry, 1994 Update. American Thoracic Society. *Am J Respir Crit Care Med* 1995; 152:1107-36.
30. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol* 1989; 129:687-702.
31. Marossy AE, Strachan DP, Rudnicka AR, Anderson HR. Childhood chest illness and the rate of decline of adult lung function between ages 35 and 45 years. *Am J Respir Crit Care Med* 2007; 175:355-9.
32. Dijkstra A, Howard TD, Vonk JM, Ampleford EJ, Lange LA, Bleecker ER, et al. Estrogen receptor 1 polymorphisms are associated with airway hyperresponsiveness and lung function decline, particularly in female subjects with asthma. *J Allergy Clin Immunol* 2006; 117:604-11.

33. Li X, Howard TD, Moore WC, Ampleford EJ, Li H, Busse WW, et al. Importance of hedgehog interacting protein and other lung function genes in asthma. *J Allergy Clin Immunol* 2011; 127:1457-65.
34. Malhotra A, Peiffer AP, Ryujin DT, Elsner T, Kanner RE, Leppert MF, et al. Further evidence for the role of genes on chromosome 2 and chromosome 5 in the inheritance of pulmonary function. *Am J Respir Crit Care Med* 2003; 168:556-61.
35. Ober C, Abney M, McPeck MS. The genetic dissection of complex traits in a founder population. *Am J Hum Genet* 2001; 69:1068-79.
36. Bouzigon E, Dizier MH, Krahenbuhl C, Lemaître A, Annesi-Maesano I, Betard C, et al. Clustering patterns of LOD scores for asthma-related phenotypes revealed by a genome-wide screen in 295 French EGEA families. *Hum Mol Genet* 2004; 13:3103-13.
37. Barton SJ, Koppelman GH, Vonk JM, Browning CA, Nolte IM, Stewart CE, et al. PLAUR polymorphisms are associated with asthma, PLAUR levels, and lung function decline. *J Allergy Clin Immunol* 2009; 123:1391-400 e17.
38. Xu X, Fang Z, Wang B, Chen C, Guang W, Jin Y, et al. A genomewide search for quantitative-trait loci underlying asthma. *Am J Hum Genet* 2001; 69:1271-7.
39. Gilliland FD, Gauderman WJ, Vora H, Rappaport E, Dubeau L. Effects of glutathione-S-transferase M1, T1, and P1 on childhood lung function growth. *Am J Respir Crit Care Med* 2002; 166:710-6.
40. Flamant C, Henrion-Caude A, Boelle PY, Bremont F, Brouard J, Delaisi B, et al. Glutathione-S-transferase M1, M3, P1 and T1 polymorphisms and severity of lung disease in children with cystic fibrosis. *Pharmacogenetics* 2004; 14:295-301.
41. Romieu I, Sienra-Monge JJ, Ramirez-Aguilar M, Moreno-Macias H, Reyes-Ruiz NI, Estela del Rio-Navarro B, et al. Genetic polymorphism of GSTM1 and antioxidant supplementation influence lung function in relation to ozone exposure in asthmatic children in Mexico City. *Thorax* 2004; 59:8-10.
42. Imboden M, Downs SH, Senn O, Matyas G, Brandli O, Russi EW, et al. Glutathione S-transferase genotypes modify lung function decline in the general population: SAPALDIA cohort study. *Respir Res* 2007; 8:2.
43. Jongepier H, Boezen HM, Dijkstra A, Howard TD, Vonk JM, Koppelman GH, et al. Polymorphisms of the ADAM33 gene are associated with accelerated lung function decline in asthma. *Clin Exp Allergy* 2004; 34:757-60.
44. Simpson A, Maniatis N, Jury F, Cakebread JA, Lowe LA, Holgate ST, et al. Polymorphisms in a disintegrin and metalloprotease 33 (ADAM33) predict impaired early-life lung function. *Am J Respir Crit Care Med* 2005; 172:55-60.
45. Sadeghnejad A, Ohar JA, Zheng SL, Sterling DA, Hawkins GA, Meyers DA, et al. Adam33 polymorphisms are associated with COPD and lung function in long-term tobacco smokers. *Respir Res* 2009; 10:21.
46. van Diemen CC, Postma DS, Vonk JM, Bruinenberg M, Schouten JP, Boezen HM. A disintegrin and metalloprotease 33 polymorphisms and lung function decline in the general population. *Am J Respir Crit Care Med* 2005; 172:329-33.
47. Dijkstra A, Vonk JM, Jongepier H, Koppelman GH, Schouten JP, ten Hacken NH, et al. Lung function decline in asthma: association with inhaled corticosteroids, smoking and sex. *Thorax* 2006; 61:105-10.
48. Cornelis MC, Agrawal A, Cole JW, Hansel NN, Barnes KC, Beaty TH, et al. The Gene, Environment Association Studies consortium (GENEVA): maximizing the knowledge

- obtained from GWAS by collaboration across studies of multiple conditions. *Genet Epidemiol* 2010; 34:364-72.
49. Caporaso N, Gu F, Chatterjee N, Sheng-Chih J, Yu K, Yeager M, et al. Genome-wide and candidate gene association study of cigarette smoking behaviors. *PLoS One* 2009; 4:e4653.
50. Weedon MN, Lango H, Lindgren CM, Wallace C, Evans DM, Mangino M, et al. Genome-wide association analysis identifies 20 loci that influence adult height. *Nat Genet* 2008; 40:575-83.
51. Sovio U, Bennett AJ, Millwood IY, Molitor J, O'Reilly PF, Timpson NJ, et al. Genetic determinants of height growth assessed longitudinally from infancy to adulthood in the northern Finland birth cohort 1966. *PLoS Genet* 2009; 5:e1000409.
52. Kang SJ, Chiang CW, Palmer CD, Tayo BO, Lettre G, Butler JL, et al. Genome-wide association of anthropometric traits in African- and African-derived populations. *Hum Mol Genet* 2010; 19:2725-38.
53. Palamarchuk A, Efanov A, Nazaryan N, Santanam U, Alder H, Rassenti L, et al. 13q14 deletions in CLL involve cooperating tumor suppressors. *Blood*; 115:3916-22.
54. Hammarsund M, Corcoran MM, Wilson W, Zhu C, Einhorn S, Sangfelt O, et al. Characterization of a novel B-CLL candidate gene--DLEU7--located in the 13q14 tumor suppressor locus. *FEBS Lett* 2004; 556:75-80.
55. Rahmatpanah FB, Carstens S, Hooshmand SI, Welsh EC, Sjahputera O, Taylor KH, et al. Large-scale analysis of DNA methylation in chronic lymphocytic leukemia. *Epigenomics* 2009; 1:39-61.
56. Pekarsky Y, Zanesi N, Croce CM. Molecular basis of CLL. *Semin Cancer Biol* 2010.
57. Cooke SL, Pole JC, Chin SF, Ellis IO, Caldas C, Edwards PA. High-resolution array CGH clarifies events occurring on 8p in carcinogenesis. *BMC Cancer* 2008; 8:288.
58. Wang X, Dockery DW, Wypij D, Fay ME, Ferris BG, Jr. Pulmonary function between 6 and 18 years of age. *Pediatr Pulmonol* 1993; 15:75-88.
59. Kunzli N, Ackermann-Liebrich U, Keller R, Perruchoud AP, Schindler C. Variability of FVC and FEV1 due to technician, team, device and subject in an eight centre study: three quality control studies in SAPALDIA. Swiss Study on Air Pollution and Lung Disease in Adults. *Eur Respir J* 1995; 8:371-6.
60. Vercelli D, Martinez FD. The Faustian bargain of genetic association studies: bigger might not be better, or at least it might not be good enough. *J Allergy Clin Immunol* 2006; 117:1303-5.
61. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J Suppl* 1993; 16:5-40.

818

819 **FIGURE LEGENDS:**

820 **Figure I:** Manhattan plots of association results for decline in lung function. A) FEV1
821 decline in non-asthmatics. B) FEV1 decline in asthmatics. C) FEV1/FVC decline in non-
822 asthmatics. D) FEV1/FVC decline in asthmatics.

823

824 **Figure II:** Association of the *DLEU7* locus with decline in FEV1 in non-asthmatics. A)
825 Regional association plot, discovery phase. B) Forest plot for rs9316500. A:
826 Chromosome position (NCBI build 36.3) and recombination rate (hg18 build). The
827 sentinel SNP is represented as a diamond and r^2 for SNPs to the sentinel SNP
828 (HapMap CEU phase II). B: The size of the square of each study reflects the
829 contributing weight to the meta-analysis, details in Table EXI.

FOOTNOTES

Footnotes to Table I:

* N comprises the maximal number of subjects who contributed to at least one GWAS analysis (either decline in FEV1 or in FEV1/FVC).

†Time spacing between the first and the second spirometry assessment.

Footnote to Table II:

* MUSK refers to *TXN/MUSK/SVEP1* locus.

Footnote to Table III:

* Associations of SNPs previously associated in cross-sectional lung function in GWAS studies, (1) Framingham¹⁵, (2) CHARGE¹⁷, (3) Spirometa¹⁶, (4) Asthmatics³³ and (5) CHARGE-Spirometa¹⁸ were assessed in the discovery cohorts only if minor allele frequency (MAF) was at least 5%. SNPs tested for associations: *ADAM19*: rs2277027, rs1422795, rs6890282; *ADCY2*: rs7710510, rs6555465; *ARMC2*: rs2798641; *C10orf11*: rs11001819; *CCDC38*: rs1036429; *CDC123*: rs7068966; *CFDP1*: rs2865531; *DAAM2*: rs3008798, rs1318002, rs2395730; *FAM13A1*: rs6830970, rs2869967; *GPR126*: rs9496346, rs6570507, rs11155242, rs7753012, rs3748069, rs171891, rs263178; *HDAC4*: rs12477314; *HHIP*: rs1032295, rs1512285, rs720485, rs1828591, rs13118928, rs1512288, rs6817273; *HTR4*: rs3995090, rs1833710; *INTS12-GSTCD-NPNT*: rs3960769, rs17035917, rs17035960, rs11727735, rs10516526, rs11731417; *KCEN2*: rs9978142; *LRP1*: rs11172113; *MECOM*: rs1344555; *MFAP2*: rs2284746; *MMP15*:

853 rs2304488, rs12447804; *MTMR3*: rs17646919; *NCR3*: rs2857595; *NOTCH4*: rs206015;
854 *ONECUT1*: rs2456526; *PID1*: rs1435867, rs1358443, rs3845823; *PTCH1*: rs10512249,
855 rs576594; *RARB*: rs1529672; *SPATA9*: rs153916; *TGFB2*: rs993925; *THSD4*:
856 rs12899618; *THSD4*: rs1568010, rs1913768; *TNS1*: rs918949, rs1035672, rs929937;
857 *ZKSCAN3*: rs6903823. Non-significant associations reported in online repository.
858 † Baseline cross-sectional lung function was calculated using Quanjer formula⁶¹.
859 ‡ Proxies tested for cross-sectional association (r^2 , D'): for rs12447804 - rs2304488
860 (0.87, 1); for rs12477314 - rs4521068 (1, 1); for rs2865531 - rs12917651 (1, 1).

TABLES

Table I: Baseline characteristics of discovery and replication cohorts, by asthma status.

		%	mean \pm SD	mean \pm SD	mean \pm SD (L)	mean \pm SD	mean \pm SD (y)	mean \pm SD (mL/y)	mean \pm SD (%/y)	%
Non-asthmatics	N*	Men	Age	Height	FEV1	FEV1/FVC	Follow-up length†	annual decline FEV1	annual decline FEV1/FVC	Never smokers
Discovery (ESE-cohorts)										
EGEA	529	45.2	41.4 \pm 11.7	1.68 \pm 0.08	3.45 \pm 0.78	0.83 \pm 0.06	11.2 \pm 1.0	-28.6 \pm 25.7	-0.47 \pm 0.53	46.5
SAPALDIA	805	49.2	41.8 \pm 11.1	1.70 \pm 0.09	3.62 \pm 0.81	0.79 \pm 0.07	10.9 \pm 0.2	-34.0 \pm 28.3	-0.40 \pm 0.46	43.1
ECRHS	1343	49.7	34.1 \pm 7.1	1.70 \pm 0.10	3.81 \pm 0.83	0.83 \pm 0.06	8.9 \pm 0.9	-26.3 \pm 30.7	-0.30 \pm 0.50	40.7
Replication with in silico data										
ARIC	7156	46.3	54.5 \pm 5.6	1.69 \pm 0.09	3.01 \pm 0.75	0.75 \pm 0.07	2.9 \pm 0.2	-52.0 \pm 57.4	-0.19 \pm 0.98	40.8
FHS	3232	44.9	52.9 \pm 10.2	1.67 \pm 0.10	2.89 \pm 0.81	0.77 \pm 0.08	10.5 \pm 3.6	-24.9 \pm 23.9	-0.33 \pm 0.57	36.1
B58C	470	48.7	35.0 \pm 0.2	1.70 \pm 0.09	3.68 \pm 0.73	0.81 \pm 0.06	10.1 \pm 0.5	-34.9 \pm 31.4	-0.21 \pm 0.67	28.5
Asthmatics										
Discovery (ESE-cohorts)										
EGEA	330	50.6	38.5 \pm 12.5	1.70 \pm 0.09	3.26 \pm 0.91	0.77 \pm 0.11	11.6 \pm 1.0	-27.6 \pm 39.4	-0.44 \pm 0.68	44.6
SAPALDIA	540	46.5	40.2 \pm 11.3	1.69 \pm 0.09	3.36 \pm 0.89	0.76 \pm 0.95	10.9 \pm 0.3	-35.5 \pm 33.9	-0.45 \pm 0.54	42.4
ECRHS	571	42.7	33.9 \pm 7.3	1.69 \pm 0.10	3.43 \pm 0.81	0.78 \pm 0.09	8.8 \pm 0.7	-26.7 \pm 42.6	-0.20 \pm 0.60	42.5
Replication with in silico data										
ARIC	325	50.2	54.2 \pm 5.7	1.69 \pm 0.10	2.73 \pm 0.87	0.68 \pm 0.10	2.9 \pm 0.2	-43.9 \pm 77.2	-0.037 \pm 1.25	41.9
FHS	346	41.3	50.1 \pm 10.3	1.68 \pm 0.09	2.72 \pm 0.84	0.73 \pm 0.09	10.2 \pm 3.8	-29.8 \pm 23.7	-0.38 \pm 0.51	36.1
B58C	231	44.2	35.0 \pm 0.2	1.69 \pm 0.10	3.45 \pm 0.75	0.78 \pm 0.08	10.3 \pm 0.5	-34.4 \pm 37.6	-0.17 \pm 0.89	37.2
Dutch Asthma	258	60.9	35.1 \pm 7.6	1.75 \pm 0.09	3.03 \pm 0.95	0.65 \pm 0.13	14.6 \pm 7.2	-22.8 \pm 47.0	-0.14 \pm 0.89	40.7

Table II: Association of (lead) SNPs subjected to replication with A) decline in FEV1 and B) decline in FEV1/FVC; stratified by asthma status.

A - decline in FEV1					Discovery phase				Replication phase			
dbSNP ID	chr	position (build 36.3)	gene nearby	Maximal frequency of coding allele	Estimate of joint analysis	P for joint analysis	P for heterogeneity between studies	P for heterogeneity between asthmatics and non- asthmatics	Estimate of joint analysis in replication cohorts	P for joint analysis	P for heterogeneity between studies	P for heterogeneity between asthmatics and non-asthmatics
Non-Asthmatics												
rs1889321	9	112340656	MUSK*	0.287	-0.150	6.95E-07	0.814	0.0187	-0.011	0.480	0.713	0.053
rs9316500	13	49992115	DLEU7	0.336	0.135	4.81E-06	0.613	0.0255	0.033	0.026	0.124	0.075
rs6785065	3	149016533	ZIC1	0.274	-0.136	0.00001	0.234	0.1700	-0.006	0.686	0.525	0.55
rs278037	13	29322627	UBL3	0.178	-0.151	0.00002	0.364	0.0058	-0.006	0.734	0.231	0.50
rs7641198	3	117396577	LSAMP	0.147	0.164	0.00003	0.669	0.1997	-0.002	0.939	0.690	0.15
rs421847	21	19269950	PRSS7	0.281	0.128	0.00003	0.831	0.0350	-0.016	0.310	0.247	0.86
rs496809	18	74857661	SALL3	0.078	-0.236	0.00004	0.412	0.0041	0.022	0.443	0.373	0.60
rs10933964	3	110021881	TRAT1	0.499	-0.117	0.00006	0.345	0.0022	-0.015	0.265	0.869	0.041
Asthmatics												
rs10808265	7	131840229	PLXNA4B	0.484	-0.175	1.66E-06	0.844	0.0020	0.069	0.105	0.258	0.16
rs1902618	15	58951491	RORA	0.234	-0.220	1.72E-06	0.449	0.0043	0.029	0.590	0.777	0.58
rs3843306	1	91060718	BARHL2	0.460	0.176	5.11E-06	0.042	8.33E-06	0.047	0.270	0.883	0.24
rs7006290	8	41734295	ANK1	0.319	0.185	5.19E-06	0.058	0.0003	0.038	0.456	0.574	0.45
rs12436689	14	84723772	FLRT2	0.244	-0.212	6.87E-06	0.420	0.0010	-0.017	0.759	0.051	0.95
rs12615721	2	81710037	CTNNA2	0.104	-0.303	7.65E-06	0.853	0.0020	-0.127	0.129	0.824	0.08
rs10516809	4	89640109	HERC5	0.101	0.306	8.67E-06	0.790	3.60E-05	-0.060	0.446	0.200	0.41
rs16856186	1	203944749	SLC45A3	0.098	0.268	8.92E-06	0.510	0.0034	-0.079	0.350	0.094	0.46
rs158536	20	52148709	BCAS1	0.408	0.162	0.00002	0.948	0.0001	0.075	0.100	0.917	0.09
rs477725	19	42066106	ZNF345	0.158	0.223	0.00003	0.821	0.0031	-0.069	0.273	0.255	0.14
rs9662589	1	230344234	DISC1	0.221	0.188	0.00005	0.868	0.0002	-0.020	0.706	0.153	0.76


rs777433	2	128084705	LIMS2	0.407	0.151	0.00010	0.811	0.1223	-0.018	0.691	0.564	0.52
B - decline in FEV1/FVC					Discovery phase				Replication phase			
dbSNP ID	chr	position (build 36.3)	gene nearby	Maximal frequency of coding allele	Estimate of joint analysis	P for joint analysis	P for heterogeneity between studies	P for heterogeneity between asthmatics and non- asthmatics	estimate of joint analysis in replication cohorts	P for joint analysis	P for heterogeneity between studies	P for heterogeneity between asthmatics and non-asthmatics
Non-Asthmatics												
rs2658782	11	92806379	FLJ25393	0.166	0.186	4.33E-06	0.362	0.0041	0.031	0.135	0.242	0.91
rs1867982	10	73197053	C10orf54	0.109	0.202	5.56E-06	0.839	0.0034	-0.008	0.745	0.412	0.24
rs12712969	2	46185673	PRKCE	0.268	-0.147	7.08E-06	0.687	0.0116	0.012	0.448	0.916	0.76
rs10187654	2	234478798	TRPM8	0.205	0.151	8.87E-06	0.797	0.0049	-0.015	0.382	0.676	0.15
rs356642	2	100903870	NPAS2	0.189	0.158	9.79E-06	0.162	0.0014	-0.010	0.565	0.282	0.28
rs890515	8	67534388	ADHFE1	0.497	0.119	0.00001	0.580	0.0257	0.003	0.847	0.443	0.58
rs10738890	9	32448081	DDX58	0.391	-0.118	0.00003	0.832	0.5847	-0.009	0.567	0.032	0.73
Asthmatics												
rs4831760	8	15576956	TUSC3	0.326	0.222	5.27E-08	0.066	7.74E-08	0.011	0.799	0.541	0.73
rs7144584	14	63345565	SYNE2	0.116	-0.318	5.62E-07	0.616	0.0010	0.089	0.272	0.752	0.43
rs1148186	10	28657641	MPP7	0.194	0.219	7.28E-06	0.760	0.0035	-0.033	0.602	0.967	0.60

Table III : Association* of SNPs previously identified in GWAS on cross-sectional lung function with percent predicted lung function at baseline, as well as percent change and annual decline in lung function for A) FEV1 and B) FEV1/FVC in ESE-discovery cohorts by asthma status.


A						Non-Asthmatics			Asthmatics		
						FEV1 percent predicted	FEV1 percent change	FEV1 decline [%/y]	FEV1 percent predicted	FEV1 percent change	FEV1 decline [%/y]
dbSNP ID	chr	position (build 36.3)	refs	gene nearby	Maximal frequency of coding allele	P-value	P-value	P-value	P-value	P-value	P-value
rs1435867	2	229219173	2	PID1	0.065	0.021	0.845	0.418	0.824	0.321	0.377
rs17035917	4	106740191	2,3	INTS12-GSTCD-NPNT	0.071	0.006	0.135	0.077	0.073	0.056	0.061
rs17035960	4	106751295	2,3	INTS12-GSTCD-NPNT	0.071	0.004	0.093	0.054	0.067	0.056	0.063
rs11727735	4	106851319	2,3	INTS12-GSTCD-NPNT	0.076	2.14E-04	0.361	0.198	0.057	0.114	0.074
rs10516526	4	106908353	2,3	INTS12-GSTCD-NPNT	0.072	1.96E-04	0.327	0.177	0.062	0.120	0.078
rs11731417	4	106965461	2,3	INTS12-GSTCD-NPNT	0.073	1.32E-04	0.335	0.177	0.048	0.146	0.090
rs1032295	4	145654034	2	HHIP	0.397	0.173	0.096	0.306	0.274	0.042	0.033
rs1512285	4	145670409		HHIP	0.462	0.032	0.029	0.141	0.152	0.033	0.024
rs720485	4	145682038	2,3	HHIP	0.391	0.159	0.510	0.786	0.943	0.044	0.058
rs1512288	4	145710731	2,3	HHIP	0.401	0.188	0.533	0.813	0.781	0.046	0.057
rs6817273	4	145711453	2,3	HHIP	0.400	0.179	0.535	0.816	0.866	0.046	0.057
rs3008798	6	39887840	3	DAAM2	0.464	0.326	0.960	0.850	0.755	0.009	0.017
rs1318002	6	39892112	3	DAAM2	0.480	0.649	0.725	0.902	0.782	0.015	0.023
rs2395730	6	39892343	3	DAAM2	0.442	0.522	0.716	0.513	0.619	0.003	0.007
rs12899618	15	69432174	3	THSD4	0.158	0.003	0.003	0.014	0.424	0.137	0.131
rs1913768	15	69436598	3	THSD4	0.159	0.003	0.002	0.011	0.393	0.162	0.152
rs1568010	15	69455566	4	THSD4	0.372	0.535	0.042	0.067	0.413	0.241	0.111
rs2304488	16	56631711	1	MMP15	0.186	0.033	0.101	0.147	0.112	0.344	0.506
rs12447804†	16	56632783	5	MMP15	0.179	0.033	0.111	0.161	0.112	0.382	0.482

B						Non-Asthmatics			Asthmatics		
						FEV1 percent predicted	FEV1 percent change	FEV1 decline [%/y]	FEV1 percent predicted	FEV1 percent change	FEV1 decline [%/y]
dbSNP ID	chr	position (build 36.3)	refs	gene nearby	Maximal frequency of coding allele	P-value	P-value	P-value	P-value	P-value	P-value
rs918949	2	218382942	2,3	TNS1	0.384	0.010	0.133	0.089	0.076	0.241	0.256
rs1035672	2	218383444	2,3	TNS1	0.384	0.010	0.133	0.089	0.093	0.243	0.258
rs929937	2	218417460	2,4	TNS1	0.386	0.623	0.017	0.016	0.004	0.915	0.888
rs3845823	2	229611365	4	PID1	0.432	0.039	0.963	0.852	0.393	0.997	0.987
rs12477314†	2	239542085	5	HDAC4	0.215	0.023	0.727	0.655	0.125	0.361	0.278
rs1529672	3	25495586	5	RARB	0.159	0.012	0.329	0.337	0.605	0.716	0.860
rs1828591	4	145700230	2,3	HHIP	0.394	0.031	0.470	0.345	0.254	0.138	0.139
rs13118928	4	145705839	2,3	HHIP	0.393	0.043	0.500	0.371	0.271	0.132	0.132
rs3995090	5	147826008	2,3	HTR4	0.394	0.011	0.785	0.699	0.029	0.649	0.456
rs2395730	6	39892343	3	DAAM2	0.442	0.277	0.554	0.685	0.979	0.036	0.039
rs2798641	6	109374743	5	ARMC2	0.209	0.315	0.444	0.530	0.006	0.188	0.158
rs9496346	6	142711031	2	GPR126	0.316	0.053	0.378	0.368	0.098	0.777	0.788
rs6570507	6	142721265	2	GPR126	0.314	0.035	0.356	0.342	0.080	0.804	0.821
rs11155242	6	142733242	2	GPR126	0.210	0.008	0.785	0.670	0.268	0.857	0.807
rs7753012	6	142787576	2	GPR126	0.337	0.051	0.477	0.487	0.065	0.566	0.637
rs3748069	6	142809326	2	GPR126	0.319	0.043	0.407	0.401	0.134	0.604	0.628
rs171891	6	142892305	2,4	GPR126	0.198	0.013	0.884	0.741	0.129	0.830	0.815
rs10512249	9	97296130	2	PTCH1	0.089	0.435	0.922	0.999	0.807	0.032	0.028
rs11172113	12	55813550	5	LRP1	0.384	0.005	0.602	0.530	0.809	0.114	0.125
rs1036429	12	94795559	5	CCDC38	0.217	0.765	0.322	0.356	0.295	0.047	0.031
rs2456526	15	50876734	1	ONECUT1	0.136	0.011	0.524	0.500	0.451	0.230	0.250
rs12899618	15	69432174	3	THSD4	0.158	3.25E-04	0.253	0.390	0.328	0.596	0.668
rs1913768	15	69436598	3,4	THSD4	0.159	4.78E-04	0.221	0.344	0.365	0.617	0.695
rs2304488	16	56631711	1	MMP15	0.186	0.121	0.002	0.002	0.085	0.760	0.515
rs12447804†	16	56632783	5	MMP15	0.179	0.121	0.003	0.003	0.085	0.719	0.487
rs2865531†	16	73947817	5	CFDP1	0.428	0.035	0.621	0.736	0.377	0.840	0.603


Figure 1




A - FEV1 decline in non-asthmatics



B - FEV1 decline in asthmatics



C - FEV1/FVC decline in non-asthmatics



D - FEV1/FVC decline in asthmatics

Figure 2

